



## WR Methodological Module

### Estimation of baseline carbon stock changes and greenhouse gas emissions from Wetland Restoration with hydrologic management including projected wetland loss in the baseline scenario (BL-WR-HM-WL)

#### I. SCOPE, APPLICABILITY AND PARAMETERS

##### Scope

This module allows for estimating carbon stock changes and GHG emissions related to Wetland Restoration (WR) in the baseline case when the baseline includes the projected reduction of total wetland project area due to wetland loss that would occur in the baseline over a 40-year Crediting Period, and the project activity will include hydrologic management. Examples of eligible hydrologic management project activities include:

- a. Diversion of river water (e.g., Mississippi River or other) into wetlands;
- b. Introduction of nonpoint source runoff (e.g., agricultural, stormwater) into wetlands;
- c. Discharge of treated municipal effluent into wetlands (e.g., wetland assimilation).
- d. Outfall management to maximize sheet flow and minimize impounded or stagnant conditions.

Implementing a wetland restoration activity not only enhances vegetative productivity and carbon sequestration, but also may prevent further wetland loss. This module accounts for baseline sequestration that would decrease over time as wetlands degrade and convert into open water. This module ~~also conservatively does not~~ account for the fate and transport of carbon during wetland loss that may result in emissions due to oxidation of soil organic carbon.

##### Applicability

The module is applicable for estimating baseline carbon stock changes and GHG emissions related to wetland restoration (WR) through hydrologic management, or through the combination of hydrologic management with assisted natural regeneration, seeding, or tree planting. The following conditions must be met to apply this module.

- Project activities meet the applicability conditions in the **WR-MF** listed under All Activity Types, and Wetland Restoration with Hydrologic Management.
- Project activities that increase emissions beyond the baseline scenario must be accounted for.

All WR activities involving hydrologic management shall occur in compliance with all applicable local, state and federal environmental regulations. The Project Proponent shall provide attestations and/or evidence (e.g. permits) of environmental compliance to ACR at the time of GHG Project Plan submission, and to the validation/verification body at the time of validation. Any changes to the project's environmental compliance status shall be reported to ACR.

## Parameters

This module provides procedures to determine the following parameters:

Parameter	SI Unit	Description
$\Delta C_{\text{bsl,WR-HM-WL}}$	t CO <sub>2</sub> -e	Cumulative total carbon stock changes and greenhouse gas emissions for the baseline scenario including projected wetland loss that would result in a reduction of the wetland project area, when the project activity will include hydrologic management
<u><math>\Delta C_{\text{TREE BSL loss}}</math></u>	<u>t CO<sub>2</sub>-e</u>	<u>Cumulative total carbon stock changes of living tree biomass for the baseline scenario including wetland loss that would result in a reduction of the wetland project area and a loss of sequestration capacity in the absence of project implementation</u>
<u><math>\Delta C_{\text{SOC BSL loss}}</math></u>	<u>t CO<sub>2</sub>-e</u>	<u>Cumulative total carbon stock changes of soils for the baseline scenario including wetland loss that would result in a reduction of the wetland project area and a loss of sequestration capacity in the absence of project implementation</u>
<u><math>\Delta \text{GHG}_E \text{ BSL loss}</math></u>	<u>t CO<sub>2</sub>-e</u>	<u>Rate of GHG emissions from the project area with BAU practices including wetland loss that would result in a reduction of the wetland project area and GHG emissions resulting from decomposition of organic matter during wetland degradation</u>

## II. PROCEDURE

This module proceeds in five steps:

- Step 1: Identification of baseline scenario
- Step 2: Project boundary

- Step 3: Baseline stratification
- Step 4: Baseline net GHG removals for fixed baselines
- Step 5: Monitoring requirements for baseline renewal

### **Step 1. Identification of the baseline scenario**

Baseline determination is defined from Paragraph 22 of the CDM A/R Modalities and Procedures as “existing or historical, as applicable, changes in carbon stocks in the carbon pools within the project boundary” where the land would remain degraded in the absence of the project activity. Project Proponent must demonstrate that the land would remain degraded in the absence of the project activity by applying the tool **T-DEG**. This may be accomplished by using multiple sources of data, such as from peer-reviewed literature, archives, maps or satellite images of the land use/cover prior to project activity, field surveys, governmental reports, expert judgment<sup>1</sup> and interviews with land owners or professionals affiliated with wetland management of the area. Project Proponent must demonstrate that the candidate baseline scenario does not alter historical wetland patterns by analyzing the historical and existing wetland over the most recent 10-year period prior to the project start date, or longer if necessary, to demonstrate baseline historical wetland patterns.

### **Step 2. Project boundary**

The project boundary geographically delineates the WR project activity under the control of the Project Proponent (PP) as defined in the **WR-MF**. It shall be demonstrated that each discrete parcel of land to be included in the boundary is eligible for a WR ACR project activity.

The pools that will be included or excluded from accounting are provided in **WR-MF**. **WR-MF** shall be followed in determining the GHG assessment boundary, along with the guidance in the *ACR Forest Carbon Project Standard*, Chapter 2.

Hydrologic management has the potential to increase GHG emissions therefore baseline and project emissions must be accounted for. Emissions may be estimated based on site/project specific data, an acceptable proxy, reference sample plots or field monitoring of similar sites, peer-reviewed literature, approved local or national parameters, or the most recent default emission factors provided by IPCC (e.g. IPCC 1997, 2003, 2006). Project Proponent using extrapolated values must make conservative estimates to determine the baseline and proposed project activity GHG emissions. Sources deemed significant and selected for accounting in the baseline scenario shall also be accounted for in the project scenario.

Exclusion of carbon pools and emission sources is allowed subject to considerations of conservativeness and significance testing. This may be accomplished by using multiple sources

---

<sup>1</sup> Justification should be supplied for all values derived from expert judgment.

of data, such as from peer-reviewed literature, field surveys, governmental reports, and expert judgment<sup>2</sup>. Pools or sources may always be excluded if conservative, i.e. exclusion will tend to underestimate net GHG emission reductions/removal enhancements. Pools or sources can be neglected (i.e., counted as zero) if application of the tool **T-SIG** indicates that the source is insignificant, provided that all sources, sinks and pools determined to be insignificant and excluded from accounting represent less than 3% of the *ex ante* calculation of emission reductions/removal enhancements (per *ACR Forest Carbon Project Standard*). If monitoring of baseline and project emissions determines that an emission source(s) initially included in the GHG assessment boundary is insignificant using the tool **T-SIG**, monitoring may cease.

### Step 3. Baseline stratification

Stratification is a standard statistical procedure to decrease overall variability of carbon stock estimates by grouping data taken from environments with similar characteristics. When estimating baseline carbon stocks, several strata can be assessed, including but not limited to:

- a. Management regime
- b. Vegetation type and species
- c. Age class
- d. Trend in land loss conversion
- e. Water quality (e.g. salinity, nutrient inputs, distance from source, etc.)
- f. Hydrology
- g. Elevation and subsidence rates
- h. Site index and anticipated growth rates
- i. Areas prone towards wetland loss

If the project activity area is not homogeneous, stratification should be carried out to improve the accuracy and precision of carbon stock estimates. Different stratifications may be required for the baseline and project scenarios, especially if there will be a change in hydrology, in order to achieve optimal accuracy and precision of the estimates of net GHG removal by sinks. For estimation of baseline net GHG removals by sinks, or estimation of actual net GHG removals by sinks, strata should be defined based on parameters that affect GHG removals or emissions and/or that are key entry variables for the methods used to measure changes in biomass stocks.

- **For baseline net GHG removals by sinks** it will usually be sufficient to stratify according to major vegetation types since baseline removals for degraded (or degrading)

---

<sup>2</sup> Justification should be supplied for all values derived from expert judgment.

wetlands can be expected to be small in comparison to project removals;

- **For actual net GHG removals by sinks**, the stratification for *ex-ante* estimations shall be based on the project monitoring plan. The stratification for *ex-post* estimations shall be based on the actual implementation of the project monitoring plan. If natural or anthropogenic impacts (e.g., hurricanes) or other factors (e.g. altered hydrology) add variability to the growth pattern of the project area, then the *ex-post* stratification shall be revised accordingly.

Project Proponent may use remotely sensed data acquired close to the time of project commencement and/or the occurrence of natural or anthropogenic impacts for *ex-ante* and *ex-post* stratification.

#### **Step 4. Baseline net removals for fixed baselines**

The WR wetland baseline scenario is the carbon stock present immediately prior to site preparation or the most likely carbon stock in the absence of project implementation. Therefore, the baseline net GHG removals by sinks is the sum of the changes in carbon stocks in the selected carbon pools within the project boundary just prior to site preparation, or what would have occurred in the absence of the project activity.

Under the applicability conditions of this methodology:

- Changes in the carbon stock of aboveground biomass of non-tree vegetation may be conservatively assumed to be zero for all strata in the baseline scenario;
- Changes in the carbon stock of dead wood and litter/surface debris carbon pools are conservatively omitted. Therefore, the sum of the changes in the carbon stocks of dead wood and litter carbon pools is zero for all strata in the baseline scenario;
- For wetland GHG emissions, the sum of emissions will either be set as zero or quantified based on the conclusions of Step 2. Project Proponent using extrapolated values must use conservative estimates.

The baseline net GHG removals by sinks shall be estimated using the equations in this section. When applying these equations for the *ex-ante* calculation of baseline net GHG removals by sinks, Project Proponent shall provide estimates of the values of those parameters that are not available before the start of the Crediting Period and commencement of monitoring activities. Project Proponent should retain a conservative approach in making these estimates.

#### 4.1 Baseline carbon stocks<sup>3</sup>

The net carbon stock changes in the baseline are equal to the baseline living tree biomass stock plus the soil organic carbon stock that includes projected wetland loss (a reduction in wetland project area that would occur in the baseline scenario), minus the baseline greenhouse gas emissions. The baseline net GHG removals by sinks should be determined as:

$$\Delta C_{\text{bsl,WR-HM-WL}} = \Delta C_{\text{TREE\_BSL\_loss}} + \Delta C_{\text{SOC\_BSL\_loss}} - \Delta \text{GHG}_{\text{E\_BSL\_loss}} \quad (1)$$

where:

$\Delta C_{\text{bsl,WR-HM-WL}}$  Cumulative total of the carbon stock changes and greenhouse gas emissions for the baseline scenario up to time  $t$ ; t CO<sub>2</sub>-e

$\Delta C_{\text{TREE\_BSL\_loss}}$  Cumulative total of the carbon stock changes of living tree biomass for the baseline scenario including wetland loss up to time  $t$ ; t CO<sub>2</sub>-e

$\Delta C_{\text{SOC\_BSL\_loss}}$  Cumulative total of the carbon stock changes of soils for the baseline scenario including wetland loss; t CO<sub>2</sub>-e

$\Delta \text{GHG}_{\text{E\_BSL\_loss}}$  Cumulative total GHG emissions from the project area for the baseline scenario including wetland loss, and GHG emissions resulting from decomposition of organic matter during wetland loss at year  $t$ ; ton CO<sub>2</sub>-e yr<sup>-1</sup>

#### Rate of Wetland Loss

This module requires knowledge of the rate at which the project area wetlands would be lost through the project period (i.e., area of wetland loss per year) without project implementation. The baseline wetland loss in hectares per year in the project boundary must be determined before the start of the project activity, but can be modified during the course of the project if relevant data becomes available. Project Proponent must make conservative estimates.

In the ideal scenario, the factors of wetland loss are already defined and the site has specific sources of data, such as from peer-reviewed literature, archives, maps or satellite images of the land use/cover prior to project activity, field surveys, and governmental reports to provide a justified rate of wetland loss.

If no verified rate exists, the rate can be established by examining proxy areas. Examination of proxy areas may be through original data collection (field measurements and/or remote sensing analysis) or where appropriate, use of directly applicable existing data generated from credible sources.

<sup>3</sup> Stock estimates shall occur for the pools defined through the framework module WR-MF

The following criteria for applicability of proxy areas to determine wetland loss rates must be met:

- a. The main factor(s) of wetland loss in the proxy at the start of the historical reference period must be the same as those expected to cause wetland loss in the project area during the project term. The historical reference period is defined as the historical period prior to the project Start Date that serves as the source of data for defining the baseline. Examples of factors include:
  - Flood control levees/ elimination of riverine inputs
  - Impoundment
  - Reduced sediment load
  - Salinity intrusion
  - Subsidence / Relative Sea Level Rise (RSLR)
  - Lack of freshwater
  - Lack of nutrient inputs
- b. Policies, regulations, and programs adopted at the national or regional level must influence the proxy area in the same way, or have an equivalent effect at the start of the historical reference period, taking into effect the current level of enforcement and program implementation (restoration programs).
- c. Transportation networks and human infrastructure, such as navigable rivers and canals, that increase the likelihood of wetland loss and that exist historically in the proxy area must be directly comparable to those that are expected to exist within the project area during the project term.
- d. Wetland management in the proxy area will be the same as in the project area under business as usual.

#### 4.2 Baseline carbon stock changes of the living tree biomass pool during wetland loss

( $\Delta C_{TREE\ BSL\ loss}$ )

Project Proponent may conservatively quantify carbon stock changes of living tree biomass including projected wetland loss (a reduction in wetland project area) for the baseline scenario. See the module “Estimation of Carbon Stocks in Tree Biomass” (CP-TB) for calculation of carbon stock in living trees. The estimation of carbon stock changes of living trees for the baseline scenario ( $\Delta C_{TREE\ BSL}$ ) from CP-TB will be modified to ( $\Delta C_{TREE\ BSL\ loss}$ ) using the equation below. If there is more than one stratum in the baseline scenario, the outcome should be summed over

all the strata to obtain the value for the whole project.

$$\Delta C_{TREE\_BSL\_loss} = \sum_{y=1}^t (H_y/H_0) * \frac{\Delta C_{TREE\_BSL}}{t}$$

~~$\Delta C_{TREE\_BSL\_loss} = (H_y/H_0) * \Delta C_{TREE\_BSL}$~~

(2)

where:

$\Delta C_{TREE\_BSL\_loss}$  Cumulative total of the carbon stock changes of living tree biomass for the baseline scenario including wetland loss; a reduction in wetland project area that would occur in the absence of project implementation up to time t; t CO<sub>2</sub>-e

$\Delta C_{TREE\_BSL}$  Cumulative total of the carbon stock changes of living tree biomass for the baseline scenario up to time t; t CO<sub>2</sub>-e (CP-TB)

$H_{yt}$  Wetland project area at year y time t; ha

$H_0$  Wetland project area at t=0; ha

y 1, 2, 3, 4.... time to monitoring event t; years

t year of monitoring event; years

#### 4.3 Baseline carbon stock changes of the soil pool during wetland loss ( $\Delta C_{SOC\_BSL\_loss}$ )

Project Proponents may conservatively quantify carbon stock changes of the soil pool including projected wetland loss (a reduction in wetland project area) for the baseline scenario. See the module “Estimation of Carbon Stocks in the Soil Organic Carbon Pool” (CP-S) for calculation of carbon stock sequestered in soils. The estimation of carbon stock changes of the soil in the baseline scenario ( $\Delta C_{SOC\_BSL}$ ) from CP-S will be modified to ( $\Delta C_{SOC\_BSL\_loss}$ ) using the equation below. If there is more than one stratum in the baseline scenario, the outcome should be summed over all the strata to obtain the value for the whole project.

$$\Delta C_{SOC\_BSL\_loss} = \sum_{y=1}^t (H_y/H_0) * f C_{SOC\_BSL}$$

~~$\Delta C_{SOC\_BSL\_loss} = (H_y/H_0) * \Delta C_{SOC\_BSL}$~~

(32)

where:

$\Delta C_{SOC\_BSL\_loss}$  Cumulative total of the carbon stock changes of soils for the baseline scenario including wetland loss; a reduction in wetland project area that would occur in the absence of project implementation up to time t; t CO<sub>2</sub>-e

$f_{\Delta C_{SOC\_BSL}}$	Rate of increase in soil carbon stock for the baseline scenario; t CO <sub>2</sub> -e yr <sup>-1</sup> (CP-S) <del>Cumulative total of the carbon stock changes of soils for the baseline scenario up to time t; t CO<sub>2</sub>-e (CP-S)</del>
$H_y$	Wetland project area at year y; ha
$H_o$	Wetland project area at t=0; ha
$y$	1, 2, 3, 4.... time to monitoring event t; years
$t$	year of monitoring event; years

#### 4.4 Baseline emissions during wetland loss and from the decomposition of organic matter during wetland loss ( $\Delta GHG_{E\_BSL\_loss}$ )

Currently, ~~little peer-reviewed data~~ ~~insufficient knowledge~~ exists regarding the fate and transport of carbon during wetland loss. ~~Further~~ ~~Critical~~ research is needed to determine what proportion of the soil horizon, or at what rate, this material is decomposed and results in CO<sub>2</sub> and CH<sub>4</sub> emissions. If project proponents choose to quantify the projected emissions that would occur due to wetland loss in the baseline scenario, ~~and are able to sufficiently address the fate and transport of carbon during wetland loss,~~ the equations below can be applied in lieu of equation C3.5.1 in module E-E. Project Proponents may use peer-reviewed literature and must make conservative estimates. ~~Project proponents are required to~~ Project proponents must provide supporting evidence and ~~justification arguments~~ supporting the emission values used. Expert judgment and statements by professionals affiliated with wetland management may be provided as part of the supporting evidence. ~~—~~ If the rate of wetland loss is known, but the proportion or rate of soil organic carbon released as GHGs from wetland decomposition is not, SOC<sub>50cm</sub> can be conservatively set to zero:

If the annual proportion of soil organic carbon released as GHGs is known, the following equation can be used:

$$\Delta GHG_{E\_BSL\_loss} = \sum_{y=1}^t \left( (H_y/H_o) * \Delta GHG_{E\_BSL} + H_y * SOC_{50cm} * (\%CO_2 + \%CH_4 * GWP_{CH_4} + \%N_2O * GWP_{N_2O}) \right) \quad (4)$$

where:

$\Delta GHG_{E\_BSL\_loss}$	Cumulative total GHG emissions from the project area for the baseline scenario including wetland loss, and GHG emissions resulting from decomposition of organic matter during wetland loss up to time t; t CO <sub>2</sub> -e yr <sup>-1</sup>
$\Delta GHG_{E\_BSL}$	Annual total GHG emissions from the project area for the baseline scenario (equation (1) of module E-E with T=1); t CO <sub>2</sub> -e yr <sup>-1</sup>
$H_y$	Wetland project area at year y; ha

$H_0$	Wetland project area at t=0; ha
$y$	1, 2, 3, 4.... time to monitoring event t; years
$t$	year of monitoring event; years
$SOC_{50cm}$	Amount of carbon in top 50 cm of wetland soil profile; t CO <sub>2</sub> -e ha <sup>-1</sup> (CP-S)
$\%_{CO_2}$	Proportion of soil organic carbon released as CO <sub>2</sub> emissions per year (0-1); no units
$\%_{CH_4}$	Proportion of soil organic carbon released as CH <sub>4</sub> emissions per year (0-1); no units
$GWP_{CH_4}$	Global warming potential for CH <sub>4</sub> (= 21 for the first commitment period); t CO <sub>2</sub> -e (t CH <sub>4</sub> ) <sup>-1</sup>
$\%_{N_2O}$	Proportion of soil organic carbon released as N <sub>2</sub> O emissions per year (0-1); no units
$GWP_{N_2O}$	Global warming potential for N <sub>2</sub> O (= 310 for the first commitment period); t CO <sub>2</sub> -e (t N <sub>2</sub> O) <sup>-1</sup>

**Alternatively, the following equation can be used if the annual rates of emissions of GHGs during wetland loss are known:**

$$\Delta GHG_{E\_BSL\_loss} = \sum_{y=1}^t \left( (H_y/H_0) * \Delta GHG_{E\_BSL} + H_y * (f GHG_{CO_2\_loss} + f GHG_{CH_4\_loss} * GWP_{CH_4} + f GHG_{N_2O\_loss} * GWP_{N_2O}) \right) \quad (5)$$

where:

$\Delta GHG_{E\_BSL\_loss}$  Cumulative total GHG emissions from the project area for the baseline scenario including wetland loss, and GHG emissions resulting from decomposition of organic matter during wetland loss up to time t; t CO<sub>2</sub>-e yr<sup>-1</sup>

$\Delta GHG_{E\_BSL}$  Cumulative total GHG emissions from the project area for the baseline scenario (equation (1) of module E-E with T=1); ton CO<sub>2</sub>-e yr<sup>-1</sup>

$f GHG_{CO_2\_loss}$  Rate of CO<sub>2</sub> emissions during wetland loss; t C ha<sup>-1</sup> yr<sup>-1</sup>

$f GHG_{CH_4\_loss}$  Rate of CH<sub>4</sub> emissions during wetland loss; t C ha<sup>-1</sup> yr<sup>-1</sup>

$f GHG_{N_2O\_loss}$  Rate of N<sub>2</sub>O emissions during wetland loss; t C ha<sup>-1</sup> yr<sup>-1</sup>

## Step 5. Monitoring requirements for baseline renewal

A Crediting Period for a project is a predetermined length of time for which the baseline scenario is applicable. This period of time is used for carbon quantification of offsets generated relative to its baseline.

In order to renew the Crediting Periods the Project Proponent must:<sup>4</sup>

- Re-submit the GHG Project Plan in compliance with then-current GHG Program standards and criteria;
- Re-evaluate the project baseline;
- Demonstrate additionality against then-current regulations and performance standard data.
- Use GHG program-approved baseline methods, emission factors, tools, and methodologies in effect at the time of Crediting Period renewal;
- Undergo validation by an approved validation/verification body.

#### DATA AND PARAMETERS NOT MONITORED

<b>Data /parameter:</b>	<u><math>H_0</math></u>
<b>Data unit:</b>	<u>ha</u>
<b>Used in equations:</b>	<u>2, 3</u>
<b>Description:</b>	<u>Wetland project area at the start of the project (t=0)</u>
<b>Source of data:</b>	<u>Peer-reviewed literature, field surveys, governmental reports, archives, maps and satellite images of the land use/cover prior to project activity.</u>
<b>Measurement procedures (if any):</b>	<u>N/A</u>
<b>Any comment:</b>	<u>Determined at the start of project.</u>

<b>Data /parameter:</b>	<u><math>H_y</math></u>
<b>Data unit:</b>	<u>ha</u>
<b>Used in equations:</b>	<u>2, 3</u>
<b>Description:</b>	<u>Wetland project area at monitoring event t</u>
<b>Source of data:</b>	<u>Peer-reviewed literature, field surveys, governmental reports, maps and satellite images of the land use/cover after start of project activity.</u>
<b>Measurement</b>	<u>N/A</u>

<sup>4</sup> American Carbon Registry, 2010. American Carbon Registry Forest Carbon Project Standard, version 2.1. Winrock International, Little Rock, Arkansas. <http://www.americancarbonregistry.org/carbon-accounting/forest-carbon-project-standard-v2.0>. Last Accessed: June 6, 2010. 63 pages.

<b>procedures (if any):</b>	
<b>Any comment:</b>	Shall be revisited at the time of baseline renewal. Since the equations using this parameter are integrated over yearly time steps, $H_y$ for each year can be extrapolated from the project rate of wetland loss, which can be calculated as $H_y / H_0$ when $y = t$ .

<b>Data /parameter:</b>	$h$
<b>Data unit:</b>	$ha \cdot year^{-1}$
<b>Used in equations:</b>	3
<b>Description:</b>	Baseline land loss per year in the project boundary projected over the life of the project in the absence of project implementation
<b>Source of data:</b>	Peer-reviewed literature, archives, maps or satellite images of the land use/cover prior to project activity, field surveys, and governmental reports.
<b>Measurement procedures (if any):</b>	N/A
<b>Any comment:</b>	
<b>Data /parameter:</b>	$h_{i,t}$
<b>Data unit:</b>	$\% \cdot year^{-1}$
<b>Used in equations:</b>	4, 5
<b>Description:</b>	Projected annual proportion of wetland that will be lost in stratum $i$ during year $t$
<b>Source of data:</b>	Analysis of remote sensing data and/or peer-reviewed literature, archives, maps or satellite images of the land use/cover prior to project activity, field surveys, governmental reports, expert judgment for a number of proxy areas
<b>Measurement procedures (if any):</b>	N/A
<b>Any comment:</b>	Shall be revisited at the time of baseline renewal

<b>Data /parameter:</b>	$\%_{CO_2}$
<b>Data unit:</b>	(0-1) no units
<b>Used in equations:</b>	4
<b>Description:</b>	Proportion of soil organic carbon released as $CO_2$ emissions per year

<b>Source of data:</b>	<a href="#">Peer-reviewed literature and/or independent field surveys</a>
<b>Measurement procedures (if any):</b>	<a href="#">N/A</a>
<b>Any comment:</b>	<a href="#">The use of this parameter must be scientifically justified by the Project Proponents and accepted by verifiers.</a>

<b>Data /parameter:</b>	<a href="#">%<sub>CH4</sub></a>
<b>Data unit:</b>	<a href="#">(0-1) no units</a>
<b>Used in equations:</b>	<a href="#">4</a>
<b>Description:</b>	<a href="#">Proportion of soil organic carbon released as CH<sub>4</sub> emissions per year</a>
<b>Source of data:</b>	<a href="#">Peer-reviewed literature and/or independent field surveys</a>
<b>Measurement procedures (if any):</b>	<a href="#">N/A</a>
<b>Any comment:</b>	<a href="#">The use of this parameter must be scientifically justified by the Project Proponents and accepted by verifiers.</a>

<b>Data /parameter:</b>	<a href="#">%<sub>N2O</sub></a>
<b>Data unit:</b>	<a href="#">(0-1) no units</a>
<b>Used in equations:</b>	<a href="#">4</a>
<b>Description:</b>	<a href="#">Proportion of soil organic carbon released as N<sub>2</sub>O emissions per year</a>
<b>Source of data:</b>	<a href="#">Peer-reviewed literature and/or independent field surveys</a>
<b>Measurement procedures (if any):</b>	<a href="#">N/A</a>
<b>Any comment:</b>	<a href="#">The use of this parameter must be scientifically justified by the Project Proponents and accepted by verifiers.</a>

<b>Data /parameter:</b>	<a href="#">f GHG<sub>CO2 loss</sub></a>
<b>Data unit:</b>	<a href="#">t C ha<sup>-1</sup> yr<sup>-1</sup></a>

<b>Used in equations:</b>	<u>5</u>
<b>Description:</b>	<u>Rate of CO<sub>2</sub> emissions during wetland loss</u>
<b>Source of data:</b>	<u>Peer-reviewed literature and/or independent field surveys</u>
<b>Measurement procedures (if any):</b>	<u>N/A</u>
<b>Any comment:</b>	<u>The use of this parameter must be scientifically justified by the Project Proponents and accepted by verifiers.</u>

<b>Data /parameter:</b>	<u><math>f_{GHG_{CH_4 \text{ loss}}}</math></u>
<b>Data unit:</b>	<u>t C ha<sup>-1</sup> yr<sup>-1</sup></u>
<b>Used in equations:</b>	<u>5</u>
<b>Description:</b>	<u>Rate of CH<sub>4</sub> emissions during wetland loss</u>
<b>Source of data:</b>	<u>Peer-reviewed literature and/or independent field surveys</u>
<b>Measurement procedures (if any):</b>	<u>N/A</u>
<b>Any comment:</b>	<u>The use of this parameter must be scientifically justified by the Project Proponents and accepted by verifiers.</u>

--	--

<b>Data /parameter:</b>	<u><math>f_{GHG_{N_2O \text{ loss}}}</math></u>
<b>Data unit:</b>	<u>t C ha<sup>-1</sup> yr<sup>-1</sup></u>
<b>Used in equations:</b>	<u>5</u>
<b>Description:</b>	<u>Rate of N<sub>2</sub>O emissions during wetland loss</u>
<b>Source of data:</b>	<u>Peer-reviewed literature and/or independent field surveys</u>
<b>Measurement procedures (if any):</b>	<u>N/A</u>
<b>Any comment:</b>	<u>The use of this parameter must be scientifically justified by the Project Proponents and accepted by verifiers.</u>

--	--

## PARAMETERS ORIGINATING IN OTHER MODULES

<b>Data /parameter:</b>	$\Delta C_{TREE\_BSL}$
<b>Data unit:</b>	t CO <sub>2</sub> -e
<b>Used in equations:</b>	<a href="#">12</a>
<b>Description:</b>	Cumulative total of the carbon stock changes of living tree biomass in the baseline scenario <a href="#">up to time t</a>
<b>Module parameter originates in:</b>	CP-TB
<b>Any comment:</b>	

<b>Data /parameter:</b>	$\Delta GHG_{E\_BSL}$
<b>Data unit:</b>	t CO <sub>2</sub> -e <a href="#">yr<sup>-1</sup></a>
<b>Used in equations:</b>	1
<b>Description:</b>	<a href="#">Annual total GHG emissions from the project area for the baseline scenario</a> <del>Cumulative total of the change in carbon emissions of the baseline scenario</del>
<b>Module parameter originates in:</b>	E-E
<b>Any comment:</b>	<a href="#">Use equation (1) of module E-E with T set to 1 (T=1)</a>

<b>Data /parameter:</b>	$f_{C_{SOC\_BSL}} \Delta C_{SOC\_BSL}$
<b>Data unit:</b>	t CO <sub>2</sub> -e <a href="#">yr<sup>-1</sup></a>
<b>Used in equations:</b>	<a href="#">2, 43</a>
<b>Description:</b>	<a href="#">Rate of increase in soil carbon stock for the baseline scenario</a> <del>Cumulative total of the carbon stock changes of soils for the baseline scenario</del>
<b>Module parameter originates in:</b>	CP-S
<b>Any comment:</b>	

<b><u>Data /parameter:</u></b>	<u>SOC<sub>50cm</sub></u>
<b><u>Data unit:</u></b>	<u>t CO2-e ha<sup>-1</sup></u>
<b><u>Used in equations:</u></b>	<u>4</u>
<b><u>Description:</u></b>	<u>Amount of carbon in top 50 cm of wetland soil profile</u>
<b><u>Module parameter originates in:</u></b>	<u>CP-S</u>
<b><u>Any comment:</u></b>	